

On the strange quark mass with improved staggered quarks

J. Hein^a, C. Davies^b, G. P. Lepage^c, Q. Mason^c and H. Trottier^d (HPQCD and UKQCD collaboration)

^aSchool of Physics, The University of Edinburgh, Edinburgh EH9 3JZ, Scotland, UK

^bDepartment of Physics & Astronomy, The University of Glasgow, Glasgow G12 8QQ, Scotland, UK

^cNewman Laboratory of Nuclear Studies, Cornell University, Ithaca, NY 14850, USA

^dPhysics Department, Simon Fraser University, Burnaby, B.C., Canada V5A 1S6

We present results on the sum of the masses of light and strange quark using improved staggered quarks. Our calculation uses 2+1 flavours of dynamical quarks. The effects of the dynamical quarks are clearly visible.

1. INTRODUCTION

1.1. Motivation

Simulations of hadronic matter with realistic sea quark effects form a long standing problem in lattice gauge theory. To achieve this goal on the upcoming generation of computers, improved staggered quarks are most promising. Compared to fermion formulations based on the Wilson action, staggered quarks have fewer variables and are faster to simulate for small quark masses.

Naïve staggered quarks suffer from large flavour changing strong interactions, which can be suppressed by the use of fat links [1]. This action can be further improved to $\mathcal{O}(\alpha_s a^2, a^4)$ [2], the most precise to date, see [3] for recent developments. These actions have proven to be successful to reduce the splittings in the pion spectrum [4] and the large renormalisations [5,6], as observed for naïve staggered quarks. Due to the advantages of improved staggered quarks it has been possible to generate data sets with 2 + 1 flavours of dynamical sea quarks with a low m_π/m_ρ -ratio [7,8]. An alternative proposal for the improvement of staggered quarks is the hypercubic action [9].

Staggered quarks have a residual “chiral” symmetry, which protects against additive mass renormalisations. The point of vanishing quark mass is exactly known beforehand. This removes the dependency on chiral extrapolations in the determination of the strange quark mass.

1.2. Simulation data

For our calculation, we use the simulation results of the MILC collaboration [7]. They use a fermionic action corrected to $\mathcal{O}(\alpha_s a^2, a^4)$ [2], which is referred to as ASQTAD. There are results for 0, 2 and 2+1 flavours of dynamical fermions. Their lightest quark mass gives a low $m_\pi/m_\rho \approx 0.37$. The simulation parameters of the configurations have been tuned to keep the lattice spacing fixed in units of the gluonic observable r_1 .

2. LATTICE SPACING

2.1. Upsilon spectrum

Υ -spectroscopy on the configurations of [7] is discussed in [10]. For the data set with $n_f = 2+1$ and the smallest m_π/m_ρ ratio one gets $a^{-1} = 1.6$ GeV. In the quenched approximation the inverse lattice spacing is substantially larger.

2.2. Kaon spectrum

In real world, the difference $m_V^2 - m_{PS}^2$ between the square of the mass of the vector and the pseudo-scalar meson is almost independent from the masses of the valence quarks. In principle this should yield a good method for determining the lattice spacing. Vector mesons are unstable however. Their decays will eventually be seen in dynamical lattice results, but currently it is unclear what effect this has on the measured masses.

In the quenched approximation, $m_V^2 - m_{PS}^2$ is much more sensitive to the valence quark masses

than the real world.. For data sets with dynamical fermions on the other hand the splittings $m_\rho^2 - m_\pi^2$, $m_{K^*}^2 - m_K^2$ and $m_\phi^2 - m_{\eta_s}^2$ differ by only $\approx 2\%$. On the $n_f = 2 + 1$ sets one gets $a^{-1} = 1.4$ GeV from $m_{K^*}^2 - m_K^2$. For the quenched data this procedure gives an inverse lattice spacing which is about 100 MeV smaller.

2.3. Gluonic scale

The gluonic observable r_1 is determined in a similar way to r_0 , but gives the point where $r^2 F(r) = 1$. Since the configurations have been generated for fixed r_1 , it is natural to use r_1 as an alternative to set the scale. MILC estimated $r_1 = 0.35$ fm, using $r_0 = 0.5$ fm. Neither of these quantities is easily related to a physical observable, however, and so there must be considerable uncertainty in their physical value. Taking $r_1 = 0.35$ fm gives $a^{-1} = 1.5$ GeV for all configurations.

3. QUARK MASS

3.1. Determination of the bare quark mass

We determine the sum of the bare masses of the light and strange quark by interpolation of the pseudo-scalar meson mass to the physical K with the ansatz

$$(am_{\text{PS}})^2 = a + b am_{\text{av}} + c(am_{\text{av}})^2. \quad (1)$$

With $m_{\text{av}} := \frac{1}{2}(m_1 + m_2)$ we denote the average of the valence masses. This is shown in figure 1 for the most chiral dynamical data set. The data points represent the π , K and η_s . The difference between a quadratic and linear interpolation is just visible in the upper right corner of the figure. At the physical K this difference is completely irrelevant.

At this point the advantage of not having additive mass renormalisations for staggered quarks pays back. No simulation is needed to determine the point of vanishing quark mass.

3.2. Renormalisation and matching

The renormalisation and matching to the $\overline{\text{MS}}$ -scheme is done in a three step procedure. We use the quark self-energy to convert from the lattice mass m_0 to the pole mass [5]. In a second step we convert to the $\overline{\text{MS}}$ -mass at the scale of the pole




Figure 1. Interpolation of the pseudoscalar mass to the Kaon

mass and finally run to the scale $\mu = 2$ GeV. At 1-loop level we get

$$\begin{aligned} m_{\overline{\text{MS}}} &= \left[1 + \alpha_s \left(b - \frac{4}{3\pi} - \frac{2}{\pi} \log(am_0) \right) \right] m_0 \\ &=: \left[1 + \alpha_s Z_0^{(2)} \right] m_0. \end{aligned} \quad (2)$$

Here $b := \delta m_0^{(2)}/m_0 - 2/\pi \log(am_0)$. This is the 1-loop contribution to the mass shift between the bare lattice mass and the pole mass with the logarithmic divergence taken out. For small values of am_0 this becomes a constant [5]. For $n_f = 2 + 1$ we obtain

$$-0.12 < Z_0^{(2)} < -0.04, \quad (3)$$

depending on the quantity used to fix the lattice spacing. It is interesting to note, for naïve staggered quarks, that is without fat links to suppress the flavour changes, $Z_0^{(2)} \approx 3$.

For α_s we use the average of α_P for the scales $aq = 1$ and 2, determined on these configs [11]. The difference between the two is included into the error. Preliminary calculations of the modified BLM scale aq^* [12] give values lying between 1 and 2. Because of the small $Z_0^{(2)}$, variations in q have only very small effects.




Figure 2. Sum of light and strange quark mass in the $\overline{\text{MS}}$ -scheme at a scale $\mu = 2$ GeV vs the m_π/m_ρ -ratio of the lightest sea quarks.

Our final result for the sum of the masses of the light and the strange quark in the $\overline{\text{MS}}$ -scheme is presented in figure 2. Different shapes of the symbols refer to the numbers of sea quarks. The different shadings of the symbols indicate the determination of the lattice spacing. The dashed line corresponds to the real world m_π/m_ρ -ratio. The error bars encompass statistical uncertainties, the uncertainties of the determination of a and the uncertainty of the scale setting of α_s . One needs to add an uncertainty for the 2-loop effects. At the present time we estimate this as $\mathcal{O}(\alpha_s^2) \approx 20\%$ for $n_f = 2 + 1$ and smaller otherwise.

In the quenched case, the results show a wide spread, depending on how a is determined. This spread is clearly reduced for $n_f = 2$ and even smaller for $n_f = 2 + 1$. There might even be a trend for a decrease with decreasing m_π/m_ρ . In table 1 we give preliminary results from the most chiral point of the corresponding data set.

4. SUMMARY

We presented the first ever calculation of the light quark masses using two flavours of dynamical

Table 1

Preliminary results on $m_l + m_s$ in $\overline{\text{MS}}$ -scheme.

n_f	$a(K^{*2} - K^2)$	$a(\Upsilon(1P) - \Upsilon(1S))$
0	94(9) MeV	71(7) MeV
2	83(11) MeV	71(10) MeV
2+1	78(14) MeV	71(13) MeV

ical light quarks and one flavour of dynamical strange. This has been possible due to the use of improved staggered quarks. Due to this improvement, the renormalisations are small and well controlled. The dynamical quarks clearly reduce the dependency on the quantity used to determine the lattice spacing. Compared to other lattice calculations and sum rule calculations, see [13] resp. [14] for review, our results favour a low value of the strange quark mass.

Acknowledgement

This work is supported by the EU IHP programme, NSF, NSERC and PPARC.

REFERENCES

1. T. Blum, et al., Phys. Rev. **D55** (1997) 1133.
2. G.P. Lepage, Phys. Rev. **D59** (1999) 074502.
3. Q. Mason, et al., these proceedings.
4. K. Orginos, D. Toussaint, R.L. Sugar, Phys. Rev. **D60** (1999) 054503.
5. J. Hein, Q. Mason, G.P. Lepage, H. Trottier, Nucl. Phys. B (Proc. Suppl.) 106 (2002) 236.
6. H. Trottier, et al., Nucl. Phys. B (Proc. Suppl.) 106 (2002) 856.
7. C. Bernard, et al., Phys. Rev. **D64** (2001), 054506.
8. D. Toussaint, et al., these proceedings.
9. A. Hasenfratz, F. Knechtli, Phys. Rev. **D64** (2001), 034504; for a recent review: A. Hasenfratz, these proceedings.
10. A. Gray, et al., these proceedings.
11. C. Davies, et al., these proceedings.
12. K. Hornbostel, G.P. Lepage, C. Morningstar, hep-ph/0208224.
13. V. Lubicz, Nucl. Phys. B (Proc. Suppl.) 94 (2001) 116; H. Wittig, these proceedings.
14. R. Gupta, K. Maltman, Int. J. Mod. Phys. A16S1B:591 (2001).